

Generalizations of generating functions for basic hypergeometric orthogonal polynomials

Howard S. Cohl[†], Roberto S. Costas-Santos[§], Philbert R. Hwang[‡], and Tanay V. Wakhare^{*}

[†] *Applied and Computational Mathematics Division, National Institute of Standards and Technology, Mission Viejo, CA 92694, USA*

URL: <http://www.nist.gov/itl/math/msg/howard-s-cohl.cfm>

E-mail: howard.cohl@nist.gov

[§] *Departamento de Física y Matemáticas, Universidad de Alcalá, c.p. 28871, Alcalá de Henares, Spain*

URL: <http://www.rscosan.com>

E-mail: rscosa@gmail.com

[‡] *Department of Computer Science, University of Maryland, College Park, MD 20742, USA*

E-mail: hwangphilbert@gmail.com

^{*} *Department of Mathematics, University of Maryland, College Park, MD 20742, USA*

E-mail: twakhare@gmail.com

Received March 10, 2021 in final form ???; Published online

doi:10.3842/JOURNAL.201*.***

Abstract. We derive generalized generating functions for basic hypergeometric orthogonal polynomials by applying connection relations with one extra free parameter to them. In particular, we generalize generating functions for the continuous q -ultraspherical/Rogers, little q -Laguerre/Wall, and q -Laguerre polynomials. Depending on what type of orthogonality these polynomials satisfy, we derive corresponding definite integrals, infinite series, bilateral infinite series, and q -integrals.

Key words: Basic hypergeometric series; Basic hypergeometric orthogonal polynomials; Generating functions; Connection coefficients; Eigenfunction expansions; Definite integrals; Infinite series; Bilateral infinite series; q -integrals.

2010 Mathematics Subject Classification: 33D45, 05A15, 33D15, 33C45, 33C20, 34L10, 30E20

1 Introduction

<Introduction>?

In the context of generalized hypergeometric orthogonal polynomials H. Cohl developed in [4, (2.1)] therein a series rearrangement technique which produces a generalization of the generating function for the Gegenbauer polynomials. We have since demonstrated that this technique is valid for a larger class of hypergeometric orthogonal polynomials. For instance, in [3] we applied this same technique to the Jacobi polynomials, and in Cohl et al. [6], we extended this technique to many generating functions for the Jacobi, Gegenbauer, Laguerre, and Wilson polynomials.

The series rearrangement technique combines a connection relation with a generating function, resulting in a series with multiple sums. The order of summations are then rearranged and the result often simplifies to produce a generalized generating function whose coefficients are given in terms of generalized or basic hypergeometric functions. This technique is especially productive when using connection relations with one extra free parameter, since the relation is most often a product of Pochhammer and q -Pochhammer symbols.

Basic hypergeometric orthogonal polynomials with more than one extra free parameter, such as the Askey-Wilson polynomials, have multi-parameter connection relations. These connection

relations are in general given by single or multiple summation expressions. For the Askey-Wilson polynomials, the connection relation with four extra free parameters is given as a basic double hypergeometric series. The fact that the four extra free parameter connection coefficient for the Askey-Wilson polynomials is given by a double sum was known to Askey and Wilson as far back as 1985 (see [9, p. 444]). When our series rearrangement technique is applied to cases with more than one extra free parameter, the resulting coefficients of the generalized generating function are rarely given in terms of a basic hypergeometric series. The more general problem of generalized generating functions with more than one extra free parameter requires the theory of multiple basic hypergeometric series and is not treated in this paper.

In this paper, we apply this technique to generalize generating functions for basic hypergeometric orthogonal polynomials in the q -analog of the Askey scheme [10, Chapter 14]. In Section 2, we give some preliminary material which is used in the remainder of the paper. In Section 3, we present generalizations of the continuous q -ultraspherical/Rogers polynomials. In Section 4, we present generalizations of the little q -Laguerre polynomials. In Section 5, we present generalizations of the q -Laguerre polynomials. In Section 6, we have also computed new definite integrals, infinite series, and Jackson integrals (hereafter q -integrals) corresponding to our generalized generating function expansions using orthogonality for the studied basic hypergeometric orthogonal polynomials.

Note that one important class of hypergeometric orthogonal polynomial generating functions which does not seem amenable to our series rearrangement technique are bilinear generating functions. The existence of an extra orthogonal polynomial in the generating function, produces multiple summation expressions via the introduction of connection relations for one or both of the polynomials with the sums being formidable to evaluate in closed form.

2 Preliminaries

(Preliminaries) Define $\mathbb{N}_0 := \{0\} \cup \mathbb{N} := \{0\} \cup \{1, 2, 3, \dots\}$. Throughout the paper, we will adopt the following notation to indicate sequential positive and negative elements, in a list of elements, namely

$$\pm a := \{a, -a\}.$$

If \pm appears in an expression, but not in a list, it is to be treated as normal. In order to obtain our derived identities, we rely on properties of the q -Pochhammer symbol (q -shifted factorial). The Pochhammer and q -Pochhammer symbols are defined for all $n \in \mathbb{N}_0$, $a \in \mathbb{C}$ such that

$$\begin{aligned} (a)_0 &:= 1, & (a)_n &:= (a)(a+1) \cdots (a+n-1), & n \in \mathbb{N}, \\ (a; q)_0 &:= 1, & (a; q)_n &:= (1-a)(1-aq) \cdots (1-aq^{n-1}), & n \in \mathbb{N}. \end{aligned} \tag{2.1} \boxed{2:1}$$

Note $(a)_b := \Gamma(a+b)/\Gamma(a)$ for all $a, b \in \mathbb{C}$, $a+b \notin -\mathbb{N}_0$. Also define

$$(a; q)_\infty := \prod_{n=0}^{\infty} (1-aq^n), \tag{2.2} \boxed{2:2}$$

where $0 < |q| < 1$, $a \in \mathbb{C}$. We will also use the common notational product conventions

$$\begin{aligned} (a_1, \dots, a_k)_b &:= (a_1)_b \cdots (a_k)_b, \\ (a_1, \dots, a_k; q)_b &:= (a_1; q)_b \cdots (a_k; q)_b, \end{aligned}$$

where $a, b \in \mathbb{C}$. We define the q -factorial as [8, (1.2.44)]

$$[0]_q! := 1, \quad [n]_q! := [1]_q [2]_q \cdots [n]_q, \quad n \in \mathbb{N},$$

where the q -number is defined as [10, (1.8.1)]

$$[z]_q := \frac{1 - q^z}{1 - q}, \quad z \in \mathbb{C},$$

with $q \in \mathbb{C}, q \neq 1$. Note that $[n]_q! = (q; q)_n / (1 - q)^n$.

The following properties for the q -Pochhammer symbol can be found in [10, (1.8.7), (1.8.10-11), (1.8.14), (1.8.19), (1.8.21-22)], namely for appropriate values of a and $k \in \mathbb{N}_0$,

$$(a^{-1}; q)_n = \frac{(-1)^n}{a^n} q^{\binom{n}{2}} (a; q^{-1})_n, \tag{2.3} \boxed{2:3}$$

$$(a; q)_{n+k} = (a; q)_k (aq^k; q)_n = (a; q)_n (aq^n; q)_k, \tag{2.4} \boxed{2:4}$$

$$(aq^n; q)_k = \frac{(a; q)_k}{(a; q)_n} (aq^k; q)_n, \tag{2.5} \text{?2:5?}$$

$$(aq^{-n}; q)_k = q^{-nk} \frac{(q/a; q)_n}{(q^{1-k}/a; q)_n} (a; q)_k, \tag{2.6} \text{?2:6?}$$

$$(a; q)_{2n} = (a, aq; q^2)_n, \tag{2.7} \text{?2:7?}$$

$$(a^2; q^2)_n = (\pm a; q)_n. \tag{2.8} \boxed{2:8}$$

Observe that by using (2.1) and (2.8), we get

$$(aq^n; q)_n = \frac{(\pm\sqrt{a}, \pm\sqrt{aq}; q)_n}{(a; q)_n}. \tag{2.9} \text{?2:9?}$$

?<negQPSqr>?

Lemma 2.1. *Let $q, \alpha, \beta \in \mathbb{C}, 0 < |q| < 1$. Then*

$$\lim_{q \uparrow 1^-} \frac{(q^\alpha; q)_\beta}{(1 - q)^\beta} = (\alpha)_\beta. \tag{2.10} \text{?2:10?}$$

Proof. See [5, Lemma 2.2]. ■

We also take advantage of the q -binomial [10, (1.11.1)] and binomial [10, (1.5.1)] theorems, $a \in \mathbb{C}, |z| < 1$, respectively $|q| < 1$,

$${}_1\phi_0 \left(\begin{matrix} a \\ - \end{matrix}; q, z \right) = \frac{(az; q)_\infty}{(z; q)_\infty},$$

where we have used (2.2), and

$${}_1F_0 \left(\begin{matrix} a \\ - \end{matrix}; q, z \right) = (1 - z)^{-a}.$$

The basic hypergeometric series, which we will often use, is defined for $|z| < 1, 0 < |q| < 1, s, r \in \mathbb{N}_0, a_l, b_j \in \mathbb{C}, l, j \in \mathbb{N}_0, 0 \leq l \leq r, 0 \leq j \leq s, b_j \notin -\mathbb{N}_0$, as [10, (1.10.1)]

$${}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, z \right) := \sum_{k=0}^{\infty} \frac{(a_1, \dots, a_r; q)_k}{(q, b_1, \dots, b_s; q)_k} \left((-1)^k q^{\binom{k}{2}} \right)^{1+s-r} z^k. \tag{2.11} \boxed{2:12}$$

Note that [10, p. 15]

$$\lim_{q \uparrow 1^-} {}_r\phi_s \left(\begin{matrix} q^{a_1}, \dots, q^{a_r} \\ q^{b_1}, \dots, q^{b_s} \end{matrix}; q, (q - 1)^{1+s-r} z \right) = {}_rF_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; z \right) := \sum_{k=0}^{\infty} \frac{(a_1, \dots, a_r)_k}{(b_1, \dots, b_s)_k} \frac{z^k}{k!}, \tag{2.12} \{?\}$$

where ${}_rF_s$ is the generalized hypergeometric series [7, Chapter 16].

Let us prove some inequalities that we will later use.

Lemma 2.2. *Let $j \in \mathbb{N}$, $k, n \in \mathbb{N}_0$, $z \in \mathbb{C}$, $\Re u > 0$, $v \geq 0$, and $0 < |q| < 1$. Then*

$$|(q^u; q)_j| \geq |(1 - |q|)[\Re u]_q|(q; q)_{j-1}|, \quad (2.13)$$

$$\left| \frac{(q^u; q)_n}{(q; q)_n} \right| \leq |[n + 1]_q^{u+1}|, \quad (2.14)$$

$$\left| \frac{(q^{v+k}; q)_n}{(q^{u+k}; q)_n} \right| \leq \frac{[n + 1]_q^{v+1}}{[\Re(u)]_q}. \quad (2.15)$$

Proof. See [5, Lemma 2.3]. ■

For a family of orthogonal polynomials $(P_n(x; \mathbf{a}))$, where \mathbf{a}, \mathbf{b} , are sets of free parameters, define $a_n(\mathbf{a})$, $c_{k,n}(\mathbf{a}; \mathbf{b})$ as follows. A generating function for these orthogonal polynomials is defined as

$$f(x, t, \mathbf{a}) = \sum_{n=0}^{\infty} a_n(\mathbf{a}) P_n(x; \mathbf{a}) t^n,$$

and a connection relation for these orthogonal polynomials is defined as

$$P_n(x; \mathbf{a}) = \sum_{k=0}^n c_{k,n}(\mathbf{a}; \mathbf{b}) P_k(x; \mathbf{b}).$$

3 The continuous q -ultraspherical/Rogers polynomials

The continuous q -ultraspherical/Rogers polynomials are defined as [10, (14.10.17)]

$$C_n(x; \beta | q) := \frac{(\beta; q)_n}{(q; q)_n} e^{in\theta} {}_2\phi_1 \left(\begin{matrix} q^{-n}, \beta \\ \beta^{-1} q^{1-n} \end{matrix}; q, q\beta^{-1} e^{-2i\theta} \right), \quad x = \cos \theta.$$

By starting with generating functions for the continuous q -ultraspherical/Rogers polynomials [10, (14.10.27–33)], we derive generalizations using the connection relation for these polynomials, namely [9, (13.3.1)]

$$C_n(x; \beta | q) = \frac{1}{1 - \gamma} \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{(1 - \gamma q^{n-2k}) \gamma^k (\beta \gamma^{-1}; q)_k (\beta; q)_{n-k}}{(q; q)_k (q\gamma; q)_{n-k}} C_{n-2k}(x; \gamma | q). \quad (3.1)$$

Theorem 3.1. *Let $x \in [-1, 1]$, $0 < |\beta|, |\gamma|, |q| < 1$, $|t\beta|(1 - |q|)^2 < 1 - |\beta|^2$. Then*

$$(te^{-i\theta}; q)_{\infty} {}_2\phi_1 \left(\begin{matrix} \beta, \beta e^{2i\theta} \\ \beta^2 \end{matrix}; q, te^{-i\theta} \right) = \sum_{n=0}^{\infty} \frac{(\beta; q)_n q^{\binom{n}{2}} (-\beta t)^n}{(\gamma, \beta^2; q)_n} C_n(x; \gamma | q) \\ \times {}_2\phi_5 \left(\begin{matrix} \beta \gamma^{-1}, \beta q^n \\ \gamma q^{n+1}, \pm \beta q^{n/2}, \pm \beta q^{(n+1)/2} \end{matrix}; q, \gamma (\beta t)^2 q^{2n+1} \right). \quad (3.2)$$

Proof. A generating function for continuous q -ultraspherical/Rogers polynomials can be found in [10, (14.10.29)]

$$(te^{-i\theta}; q)_{\infty} {}_2\phi_1 \left(\begin{matrix} \beta, \beta e^{2i\theta} \\ \beta^2 \end{matrix}; q, te^{-i\theta} \right) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-\beta t)^n}{(\beta^2; q)_n} C_n(x; \beta | q). \quad (3.3)$$

Start with (3.3), inserting (3.1), shifting the n index by $2k$, reversing the order of summation and using (2.4) through (2.11), and by noting

$$\binom{n+2k}{2} = \binom{n}{2} + 4\binom{k}{2} + (2n+1)k.$$

Define

$$\langle \mathbf{u}_R(\beta|q), f(x) \rangle := \int_{-1}^1 f(x) \frac{w_R(x; \beta|q)}{\sqrt{1-x^2}} dx. \quad (3.4) \text{ Rogersfunctional}$$

where $w_R : (-1, 1) \rightarrow [0, \infty)$ is the weight function defined by

$$w_R(x; \beta|q) := \left| \frac{(e^{2i\theta}; q)_\infty}{(\beta e^{2i\theta}; q)_\infty} \right|^2. \quad (3.5) \text{ Rogersweight}$$

This completes the proof since ([10, (14.10.18)] $|C_n(x; \beta|q)| \leq K_1[n+1]^{\sigma_1}$. **[HSC: If this completes the proof, then what is all the stuff following this sentence?]** Hence

$$|\langle \mathbf{u}, C_n(x; \beta|q) C_k(x; \beta|q) \rangle| \leq K_1^2 [n+1]^{2\sigma_1},$$

and

$$|\langle \mathbf{u}, C_n(x; \beta|q) C_n(x; \beta|q) \rangle| = \left| \frac{(\beta^2; q)_n (1-\beta)}{(q; q)_n (1-\beta q^n)} \right| \geq \left| \frac{[\Re(2b)]_q [\Re(b)]_q}{[2]_q^{b+2n+1}} \right| \geq \frac{K_2}{(1-|q|)^{2n}},$$

where $q^b = \beta$, so

$$|c_{k,n}| = \left| \frac{\langle \mathbf{u}, C_n(x; \beta|q) C_k(x; \beta|q) \rangle}{\langle \mathbf{u}, C_n(x; \beta|q) C_n(x; \beta|q) \rangle} \right| \leq \frac{K_1^2}{K_2} [n+1]^{2\sigma_1} (1-|q|)^{2n},$$

$|a_n| \leq (|t\beta|/(1-|\beta|^2))^n$. Therefore

$$\sum_{n=0}^{\infty} |a_n| \sum_{k=0}^{\lfloor n/2 \rfloor} |c_{k,n}| |C_k(x; \beta|q)| \leq \frac{K_1^3}{K_2} \sum_{n=0}^{\infty} \frac{(1-|q|)^{2n} |t|^n |\beta|^2}{(1-|\beta|^2)^n} (n+1)^{3\sigma_1+1} < \infty,$$

and the result follows. ■

Corollary 3.2. *Let $x \in [-1, 1]$, $|t| < 1$, $\beta, \gamma \in (-1, \infty) \setminus \{0, 1\}$, $0 < |q| < 1$. Then*

$$e^{xt} {}_0F_1 \left(\begin{matrix} - \\ \beta + \frac{1}{2} \end{matrix}; \frac{(x^2-1)t^2}{4} \right) = \sum_{n=0}^{\infty} \frac{(\beta)_n t^n}{(\gamma, 2\beta)_n} C_n^\gamma(x) {}_2F_3 \left(\begin{matrix} \beta - \gamma, \beta + n \\ \gamma + n + 1, \beta + \frac{n}{2}, \beta + \frac{n+1}{2} \end{matrix}; \frac{t^2}{4} \right). \quad (3.6) \text{ qto1genofgenthm2}$$

(cor4)

Proof. In (3.2), transform $\beta \mapsto q^\beta$, $\gamma \mapsto q^\gamma$, $t \mapsto (1-q)t$, and take the limit as $q \uparrow 1^-$. Using the definition of the q -exponential function [10, (1.14.2)] $E_q(z) := (-z; q)_\infty$, $\lim_{q \uparrow 1^-} E_q((1-q)z) = e^z$, and that the ${}_2\phi_1$ becomes a Kummer confluent hypergeometric functions ${}_1F_1$ with argument $-2it \sin \theta$. Representing this as a Bessel function of the first kind using [7, (10.16.5)], and then using [7, (10.2.2)], the left-hand side follows. The $q \uparrow 1^-$ limit on the right-hand side is straightforward. ■

?(theo:5)? **Theorem 3.3.** *Let $x \in [-1, 1]$, $|t|(1-|q|)^2 < 1 - |\beta|^2$, $0 < |\beta|, |\gamma|, |q| < 1$. Then*

$$\frac{1}{(te^{i\theta}; q)_\infty} {}_2\phi_1 \left(\begin{matrix} \beta, \beta e^{2i\theta} \\ \beta^2 \end{matrix}; q, te^{-i\theta} \right) = \sum_{n=0}^{\infty} \frac{(\beta; q)_n t^n}{(\gamma, \beta^2; q)_n} C_n(x; \gamma|q) {}_6\phi_5 \left(\begin{matrix} \beta\gamma^{-1}, \beta q^n, 0, 0, 0, 0 \\ \gamma q^{n+1}, \pm\beta q^{\frac{n}{2}}, \pm\beta q^{\frac{n+1}{2}} \end{matrix}; q, \gamma t^2 \right) \quad (3.7) \text{ theorem3}$$

Proof. A generating function for the continuous q -ultraspherical/Rogers polynomials can be found in [10, (14.10.28)]

$$\frac{1}{(te^{i\theta}; q)_\infty} {}_2\phi_1 \left(\begin{matrix} \beta, \beta e^{2i\theta} \\ \beta^2 \end{matrix}; q, te^{-i\theta} \right) = \sum_{n=0}^{\infty} \frac{C_n(x; \beta|q)}{(\beta^2; q)_n} t^n. \quad (3.8) \quad \boxed{\text{genfun141028}}$$

The proof follows as above by starting with (3.8), inserting (3.1), shifting the n index by $2k$, reversing the order of summation and using (2.4) through (2.11). ■

Remark 3.4. The $q \uparrow 1^{-1}$ limit of (3.7) can also be shown to be the same as (3.6), by using the transformation $x \mapsto -x$. The proof of this is the same as the proof of Corollary 3.2, except instead use the definition of the q -exponential function [10, (1.14.1)] $e_q(z) := 1/(z; q)_\infty$, $\lim_{q \uparrow 1^-} e_q((1-q)z) = e^z$. Of course, the same is true for the $q \uparrow 1^-$ limits of the original generating functions [10, (14.10.28–29)], which both are analogues of [10, (9.8.31)], and are equivalent under the transformation $x \mapsto -x$.

^{?(theo:6)?} **Theorem 3.5.** Let $x \in [-1, 1]$, $|t|(1-|q|)^2 < 1$, $0 < |\beta|, |\gamma|, |q| < 1$. Then

$$\begin{aligned} \frac{(\gamma te^{i\theta}; q)_\infty}{(te^{i\theta}; q)_\infty} {}_3\phi_2 \left(\begin{matrix} \gamma, \beta, \beta e^{2i\theta} \\ \beta^2, \gamma te^{i\theta} \end{matrix}; q, te^{-i\theta} \right) &= \sum_{n=0}^{\infty} \frac{(\beta, \gamma; q)_n t^n}{(\alpha, \beta^2; q)_n} C_n(x; \alpha|q) \\ &\times {}_6\phi_5 \left(\begin{matrix} \beta/\alpha, \beta q^n, \pm(\gamma q^n)^{\frac{1}{2}}, \pm(\gamma q^{n+1})^{\frac{1}{2}} \\ \alpha q^{n+1}, \pm\beta q^{n/2}, \pm\beta q^{(n+1)/2} \end{matrix}; q, \alpha t^2 \right). \end{aligned} \quad (3.9) \quad \boxed{\text{genFunc4C}}$$

Proof. A generating function for the continuous q -ultraspherical/Rogers polynomials can be found in [10, (14.10.33)]

$$\frac{(\gamma te^{i\theta}; q)_\infty}{(te^{i\theta}; q)_\infty} {}_3\phi_2 \left(\begin{matrix} \gamma, \beta, \beta e^{2i\theta} \\ \beta^2, \gamma te^{i\theta} \end{matrix}; q, te^{-i\theta} \right) = \sum_{n=0}^{\infty} \frac{(\gamma; q)_n}{(\beta^2; q)_n} C_n(x; \beta|q) t^n, \quad (3.10) \quad \boxed{\text{genfun4}}$$

where $\gamma \in \mathbb{C}$. Substitute (3.1) into the generating function (3.10), reverse the order of summation as above, shift the n index by $2k$, using (2.4) through (2.11), completes the proof. ■

^{?(theo:7)?} **Theorem 3.6.** Let $x \in [-1, 1]$, $(1-|q|)^2(1+|\sqrt{q}||\beta|)|t| < (1-|q||\beta|)$, $0 < |\beta|, |\gamma|, |q| < 1$. Then

$$\begin{aligned} {}_2\phi_1 \left(\begin{matrix} \pm\beta^{\frac{1}{2}} e^{i\theta} \\ -\beta \end{matrix}; q, te^{-i\theta} \right) {}_2\phi_1 \left(\begin{matrix} \pm(q\beta)^{\frac{1}{2}} e^{-i\theta} \\ -q\beta \end{matrix}; q, te^{i\theta} \right) &= \sum_{n=0}^{\infty} \frac{(\beta, \pm\beta q^{\frac{1}{2}}; q)_n t^n}{(\gamma, \beta^2, -q\beta; q)_n} C_n(x; \gamma|q) \\ &\times {}_{10}\phi_9 \left(\begin{matrix} \beta\gamma^{-1}, \beta q^n, \pm(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}}, \pm i(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm i(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \\ \gamma q^{n+1}, \pm\beta q^{n/2}, \pm\beta q^{(n+1)/2}, \pm i(\beta q^{n+1})^{\frac{1}{2}}, \pm i(\beta q^{n+2})^{\frac{1}{2}} \end{matrix}; q, \gamma t^2 \right). \end{aligned} \quad (3.11) \quad \boxed{\text{genFunthm5}}$$

Proof. A generating function for the continuous q -ultraspherical/Rogers polynomials can be found in [10, (14.10.31)]

$${}_2\phi_1 \left(\begin{matrix} \pm\beta^{\frac{1}{2}} e^{i\theta} \\ -\beta \end{matrix}; q, te^{-i\theta} \right) {}_2\phi_1 \left(\begin{matrix} \pm(q\beta)^{\frac{1}{2}} e^{-i\theta} \\ -q\beta \end{matrix}; q, te^{i\theta} \right) = \sum_{n=0}^{\infty} \frac{(\pm\beta q^{\frac{1}{2}}; q)_n}{(\beta^2, -q\beta; q)_n} C_n(x; \beta|q) t^n. \quad (3.12) \quad \boxed{\text{genfun2}}$$

We substitute (3.1) into the generating function (3.12), switch the order of the summation, shift the n index by $2k$ and using (2.4) through (2.11), produces

$$|a_n| \leq K_3 \frac{(1+|\sqrt{q}||\beta|)^n |t|^n}{(1-|q||\beta|)^n} [n+1]^{\sigma_2}.$$

Therefore the theorem holds. ■

?(theo:8)? **Theorem 3.7.** Let $x \in [-1, 1]$, $(1 - |q|)^2(1 + |\beta|\sqrt{|q|})(1 + |\beta|)|t| < (1 - |\sqrt{q}||\beta|)(1 - |\beta|^2)$, $0 < |\beta|, |\gamma|, |q| < 1$. Then

$$\begin{aligned} & {}_2\phi_1 \left(\begin{matrix} \beta^{\frac{1}{2}} e^{i\theta}, (q\beta)^{\frac{1}{2}} e^{i\theta} \\ \beta q^{\frac{1}{2}} \end{matrix}; q, te^{-i\theta} \right) {}_2\phi_1 \left(\begin{matrix} -\beta^{\frac{1}{2}} e^{-i\theta}, -(q\beta)^{\frac{1}{2}} e^{-i\theta} \\ \beta q^{\frac{1}{2}} \end{matrix}; q, te^{i\theta} \right) \\ &= \sum_{n=0}^{\infty} \frac{(\pm\beta, -\beta q^{\frac{1}{2}}; q)_n t^n}{(\gamma, \beta^2, \beta q^{\frac{1}{2}}; q)_n} C_n(x; \gamma|q) \\ &\times {}_{10}\phi_9 \left(\begin{matrix} \beta\gamma^{-1}, \beta q^n, \pm i(\beta q^n)^{\frac{1}{2}}, \pm i(\beta q^{n+1})^{\frac{1}{2}}, \pm i(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm i(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \\ \gamma q^{n+1}, \pm \beta q^{n/2}, \pm \beta q^{(n+1)/2}, \pm (\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm (\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \end{matrix}; q, \gamma t^2 \right). \end{aligned}$$

Proof. We start with the generating function for the continuous q -ultraspherical/Rogers polynomials [10, (14.10.30)]

$$\begin{aligned} & {}_2\phi_1 \left(\begin{matrix} \beta^{\frac{1}{2}} e^{i\theta}, (q\beta)^{\frac{1}{2}} e^{i\theta} \\ \beta q^{\frac{1}{2}} \end{matrix}; q, te^{-i\theta} \right) {}_2\phi_1 \left(\begin{matrix} -\beta^{\frac{1}{2}} e^{-i\theta}, -(q\beta)^{\frac{1}{2}} e^{-i\theta} \\ \beta q^{\frac{1}{2}} \end{matrix}; q, te^{i\theta} \right) \\ &= \sum_{n=0}^{\infty} \frac{(-\beta, -\beta q^{\frac{1}{2}}; q)_n}{(\beta^2, \beta q^{\frac{1}{2}}; q)_n} C_n(x; \beta|q) t^n. \end{aligned} \quad (3.13) \quad \boxed{\text{genFunc1}}$$

Using the connection relation (3.1) in (3.13), reversing the orders of the summation, shifting the n index by $2k$, and using (2.4) through (2.11), obtains the result

$$|a_n| \leq \frac{(1 + |\beta|\sqrt{|q|})^n (1 + |\beta|)^n |t|^n}{(1 - |\beta|^2)^n (1 - |\sqrt{q}||\beta|)^n}.$$

Therefore the theorem holds. ■

?(theo:9)? **Theorem 3.8.** Let $x \in [-1, 1]$, $(1 - |q|)^2(1 + |\beta|)|t| < (1 - |\sqrt{q}||\beta|)$, $0 < |\beta|, |\gamma|, |q| < 1$. Then

$$\begin{aligned} & {}_2\phi_1 \left(\begin{matrix} \beta^{\frac{1}{2}} e^{i\theta}, -(q\beta)^{\frac{1}{2}} e^{i\theta} \\ -\beta q^{\frac{1}{2}} \end{matrix}; q, te^{-i\theta} \right) {}_2\phi_1 \left(\begin{matrix} (q\beta)^{\frac{1}{2}} e^{-i\theta}, -\beta^{\frac{1}{2}} e^{-i\theta} \\ -\beta q^{\frac{1}{2}} \end{matrix}; q, te^{i\theta} \right) \\ &= \sum_{n=0}^{\infty} \frac{(\pm\beta, \beta q^{\frac{1}{2}}; q)_n t^n}{(\gamma, \beta^2, -\beta q^{\frac{1}{2}}; q)_n} C_n(x; \gamma|q) \\ &\times {}_{10}\phi_9 \left(\begin{matrix} \beta\gamma^{-1}, \beta q^n, \pm i(\beta q^n)^{\frac{1}{2}}, \pm i(\beta q^{n+1})^{\frac{1}{2}}, \pm (\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm (\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \\ \gamma q^{n+1}, \pm \beta q^{n/2}, \pm \beta q^{(n+1)/2}, \pm i(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm i(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \end{matrix}; q, \gamma t^2 \right). \end{aligned} \quad (3.14) \quad \boxed{\text{genFunc3C}}$$

Proof. A generating function for the continuous q -ultraspherical/Rogers polynomials can be found in [10, (14.10.32)]

$$\begin{aligned} & {}_2\phi_1 \left(\begin{matrix} -\beta, (\beta q)^{\frac{1}{2}} \\ -\beta q^{\frac{1}{2}} \end{matrix}; q, te^{-i\theta} \right) {}_2\phi_1 \left(\begin{matrix} (\beta q)^{\frac{1}{2}} e^{-i\theta}, -\beta^{\frac{1}{2}} e^{-i\theta} \\ -\beta q^{\frac{1}{2}} \end{matrix}; q, te^{i\theta} \right) \\ &= \sum_{n=0}^{\infty} \frac{(-\beta, \beta q^{\frac{1}{2}}; q)_n}{(\beta^2, -\beta q^{\frac{1}{2}}; q)_n} C_n(x; \beta|q) t^n. \end{aligned} \quad (3.15) \quad \boxed{\text{genfun3}}$$

Similar to the proof of (3.13), we substitute (3.1) into the generating function (3.15), switch the order of the summation, shift the n sum by $2k$, and use (2.4) through (2.11), obtaining the result

$$|a_n| \leq K_4 \frac{(1 + |\beta|)^n |t|^n}{(1 - \sqrt{q}|\beta|)^n} [n + 1]^{\sigma_3}.$$

Therefore the theorem holds. ■

4 The little q -Laguerre/Wall polynomials

The little q -Laguerre/Wall polynomials are defined as [10, (14.20.1)]

$$p_n(x; a|q) := {}_2\phi_1 \left(\begin{matrix} q^{-n}, 0 \\ aq \end{matrix}; q, qx \right) = \frac{1}{(a^{-1}q^{-n}; q)_n} {}_2\phi_0 \left(\begin{matrix} q^{-n}, x^{-1} \\ - \end{matrix}; q, \frac{x}{a} \right).$$

The connection relation for little q -Laguerre/Wall polynomials can be obtained by Exercise 1.33 in [8] and using the specialization formula which connects the little q -Laguerre/Wall polynomials with the little q -Jacobi polynomials, namely [10, p. 521] $p_n(x; a|q) = p_n(x; a, 0|q)$.

Theorem 4.1. *Let $0 < |aq|, |bq|, |q| < 1$. Then the connection relation for the little q -Laguerre/Wall polynomials is given by*

$$p_n(x; a|q) = \frac{q^{-\binom{n}{2}}}{(qa; q)_n} \sum_{j=0}^n \frac{q^{\binom{j}{2} + n(n-j)} (-a)^{n-j} (q^{n-j+1}, qb; q)_j (bq^{1+j-n}/a; q)_{n-j}}{(q; q)_j} p_j(x; b|q). \quad (4.1) \quad \boxed{\text{ConCoef3}}$$

By starting with the generating function for the little q -Laguerre/Wall polynomials [10, (14.20.11)], we derive generalizations using the connection relation for these polynomials.

Theorem 4.2. *Let $0 < |aq|, |bq|, |q| < 1$, $|t| < \min\{(1 - q)(1 - aq)/a, 1\}$. Then*

$$\frac{(t; q)_\infty}{(xt; q)_\infty} {}_0\phi_1 \left(\begin{matrix} - \\ aq \end{matrix}; q, aqxt \right) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-t)^n (bq; q)_n}{(q; q)_n (aq; q)_n} p_n(x; b|q) {}_1\phi_1 \left(\begin{matrix} a/b \\ aq^{n+1} \end{matrix}; q, bq^{n+1}t \right).$$

Proof. We start with the generating function for little q -Laguerre/Wall polynomials found in [10, (14.20.11)]

$$\frac{(t; q)_\infty}{(xt; q)_\infty} {}_0\phi_1 \left(\begin{matrix} - \\ aq \end{matrix}; q, aqxt \right) = \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{n}{2}}}{(q; q)_n} p_n(x; a|q) t^n. \quad (4.2) \quad \boxed{\text{gen1Func}}$$

Using the connection relation (4.1) in (4.2), reversing the orders of the summations, shifting the n index by j , and using (2.4) through (2.11), obtains the desired result since $|a_n| = |t|^n / (1 - q)^n$, $|c_{n,k}| \leq K_8 [n + 1]^{\sigma_6}$, and

$$|p_n(x; a|q)| \leq |a|^n [n + 1]^{\sigma_7} / (1 - |aq|)^n \leq a^n (n + 1)^{\sigma_7} / (1 - aq)^n, \quad (4.3) \quad \boxed{\text{1qlageq?}}$$

where σ_6 and σ_7 are independent of n implies

$$\sum_{n=0}^{\infty} |a_n| \sum_{k=0}^n |c_{k,n}| |p_k(x; a|q)| < \infty.$$

Therefore the theorem holds. ■

5 The q -Laguerre polynomials

repolynomials) The q -Laguerre polynomials are defined as [10, (14.21.1)]

$$L_n^{(\alpha)}(x; q) = \frac{(q^{\alpha+1}; q)_n}{(q; q)_n} {}_1\phi_1 \left(\begin{matrix} q^{-n} \\ q^{\alpha+1} \end{matrix}; q, -q^{n+\alpha+1}x \right) = \frac{1}{(q; q)_n} {}_2\phi_1 \left(\begin{matrix} q^{-n}, -x \\ 0 \end{matrix}; q, q^{n+\alpha+1} \right).$$

?<theo:12)? **Theorem 5.1.** Let $\alpha, \beta \in (-1, \infty)$, $0 < |q| < 1$. The connection relation for the q -Laguerre polynomials is given as

$$L_n^{(\alpha)}(x; q) = \frac{q^{n(\alpha-\beta)}}{(q; q)_n} \sum_{j=0}^n (-1)^{n-j} q^{\binom{n-j}{2}} (q^{n-j+1}; q)_j (q^{j-n+\beta-\alpha+1}; q)_{n-j} L_j^{(\beta)}(x; q). \quad (5.1) \quad \boxed{\text{ConCoef2}}$$

Proof. One could obtain the above result by following an analogous proof as applied to the little q -Laguerre/Wall polynomials. Nevertheless the result follows by using the relation between the little q -Laguerre/Wall and the q -Laguerre polynomials [10, p. 521]. ■

By starting with generating functions for the q -Laguerre polynomials [10, (14.21.14–16)], we derive generalizations of these generating functions using the connection relation for q -Laguerre polynomials (5.1). Note however that the generating function for the q -Laguerre polynomials [10, (14.21.13)] remains unchanged when one applies the connection relation (5.1).

?<theo:13)? **Theorem 5.2.** Let $\alpha, \beta \in (-1, \infty)$, $0 < |q| < 1$, $|t| < (1 - q^{\alpha+1})(1 - q)$. Then

$$\frac{1}{(t; q)_\infty} {}_0\phi_1 \left(\begin{matrix} - \\ q^{\alpha+1} \end{matrix}; q, -xtq^{\alpha+1} \right) = \sum_{n=0}^{\infty} \frac{(q^{\alpha-\beta}t)^n L_n^{(\beta)}(x; q)}{(q^{\alpha+1}; q)_n} {}_2\phi_1 \left(\begin{matrix} q^{\alpha-\beta}, 0 \\ q^{\alpha+n+1} \end{matrix}; q, t \right). \quad (5.2) \quad \boxed{\text{qLag1}}$$

Proof. We start with the generating function for q -Laguerre polynomials found in [10, (14.21.14)]

$$\frac{1}{(t; q)_\infty} {}_0\phi_1 \left(\begin{matrix} - \\ q^{\alpha+1} \end{matrix}; q, -xtq^{\alpha+1} \right) = \sum_{n=0}^{\infty} \frac{L_n^{(\alpha)}(x; q)}{(q^{\alpha+1}; q)_n} t^n. \quad (5.3) \quad \boxed{\text{genLFunc1}}$$

Using the connection relation (5.1) in (5.3), reversing the orders of the summations, shifting the n index by j , and using (2.4) through (2.11), obtains the desired result since $|a_n| \leq |t|^n / (1 - q^{\alpha+1})^n$, $|c_{n,k}| \leq K_9 [n+1]^{\beta-\alpha+4}$, and

$$|L_n^{(\alpha)}(x; q)| \leq [n+1]^{\sigma_8} / (1 - q)^n, \quad (5.4) \quad \boxed{\text{?qLeq?}}$$

implies

$$\sum_{n=0}^{\infty} |a_n| \sum_{k=0}^n |c_{k,n}| |L_k^{(\alpha)}(x; q)| \leq K_9 \sum_{n=0}^{\infty} \frac{|t|^n}{(1 - q^{\alpha+1})^n (1 - q)^n} (n+1)^{\sigma_8} < \infty.$$

Therefore the theorem holds. ■

?<theo:14)? **Theorem 5.3.** Let $\alpha, \beta \in (-1, \infty)$, $0 < |q| < 1$, $|t| < (1 - q^{\alpha+1})(1 - q)$. Then

$$(t; q)_\infty {}_0\phi_2 \left(\begin{matrix} - \\ q^{\alpha+1}, t \end{matrix}; q, -xtq^{\alpha+1} \right) = \sum_{n=0}^{\infty} \frac{(-tq^{\alpha-\beta})^n q^{\binom{n}{2}} L_n^{(\beta)}(x; q)}{(q^{\alpha+1}; q)_n} {}_1\phi_1 \left(\begin{matrix} q^{\alpha-\beta} \\ q^{\alpha+n+1} \end{matrix}; q, tq^n \right). \quad (5.5) \quad \boxed{\text{qLag2}}$$

Proof. We start with the generating function for the q -Laguerre polynomials found in [10, (14.21.15)]

$$(t; q)_\infty {}_0\phi_2 \left(\begin{matrix} - \\ q^{\alpha+1}, t \end{matrix}; q, -xtq^{\alpha+1} \right) = \sum_{n=0}^{\infty} \frac{(-t)^n q^{\binom{n}{2}}}{(q^{\alpha+1}; q)_n} L_n^{(\alpha)}(x; q). \quad (5.6) \text{ genLFunc2}$$

Using the connection relation (5.1) in (5.6), reversing the orders of the summations, shifting the n index by j , and using (2.4) through (2.11), obtains the desired result since, again, $|a_n| \leq |t|^n / (1 - q^{\alpha+1})^n$. ■

?(theo:15)? **Theorem 5.4.** *Let $\alpha, \beta \in (-1, \infty)$, $\gamma \in \mathbb{C}$, $0 < |q| < 1$, $|t| < 1 - q$. Then*

$$\frac{(\gamma t; q)_\infty}{(t; q)_\infty} {}_1\phi_2 \left(\begin{matrix} \gamma \\ q^{\alpha+1}, \gamma t \end{matrix}; q, -xtq^{\alpha+1} \right) = \sum_{n=0}^{\infty} \frac{(\gamma; q)_n (tq^{\alpha-\beta})^n}{(q^{\alpha+1}; q)_n} L_n^{(\beta)}(x; q) {}_2\phi_1 \left(\begin{matrix} q^{\alpha-\beta}, \gamma q^n \\ q^{\alpha+n+1} \end{matrix}; q, t \right). \quad (5.7) \text{ qLag3}$$

Proof. We start with the generating function for the q -Laguerre polynomials found in [10, (14.21.15)]

$$\frac{(\gamma t; q)_\infty}{(t; q)_\infty} {}_1\phi_2 \left(\begin{matrix} \gamma \\ q^{\alpha+1}, \gamma t \end{matrix}; q, -xtq^{\alpha+1} \right) = \sum_{n=0}^{\infty} \frac{(\gamma; q)_n}{(q^{\alpha+1}; q)_n} L_n^{(\alpha)}(x; q) t^n. \quad (5.8) \text{ genLFunc3}$$

Using the connection relation (5.1) in (5.8), reversing the orders of the summations, shifting the n index by j , and using (2.4) through (2.11), obtains the result

$$|a_n| \leq \frac{|t|^n [n+1]^{|\gamma|+1}}{\alpha+1}.$$

Therefore the theorem holds. ■

6 Definite integrals, infinite series, and q -integrals

andqintegrals Consider a sequence of orthogonal polynomials $(p_k(x; \alpha))$ (over a domain A , with positive weight $w(x; \alpha)$) associated with a linear functional \mathbf{u} , where α is a set of fixed parameters. Define s_k , $k \in \mathbb{N}_0$ by

$$s_k^2 := \int_A \{p_k(x; \alpha)\}^2 w(x; \alpha) dx.$$

In order to justify interchange between a generalized generating function via connection relation and an orthogonality relation for p_k , we show that the double sum/integral converges in the L^2 -sense with respect to the weight $w(x; \alpha)$. This requires

$$\sum_{k=0}^{\infty} d_k^2 s_k^2 < \infty, \quad (6.1) \text{ form-inve-L2}$$

where

$$d_k = \sum_{n=k}^{\infty} a_n c_{k,n}.$$

Here, a_n is the coefficient multiplying the orthogonal polynomial in the original generating function, and $c_{k,n}$ is the connection coefficient for p_k (with appropriate set of parameters).

Lemma 6.1. *Let \mathbf{u} be a classical linear functional and let $(p_n(x))$, $n \in \mathbb{N}_0$ be the sequence of orthogonal polynomials associated with \mathbf{u} . If $|p_n(x)| \leq K(n+1)^\sigma \gamma^n$, with K , σ and γ constants independent of n , then $|s_n| \leq K(n+1)^\sigma \gamma^n |s_0|$.*

Proof. See [5, Lemma 6.1]. ■

Given $|p_k(x; \boldsymbol{\alpha})| \leq K(k+1)^\sigma \gamma^k$, with K , σ and γ constants independent of k , an orthogonality relation for p_k , and $|t| < 1/\gamma$, one has $\sum_{n=0}^{\infty} |a_n| \sum_{k=0}^n |c_{k,n} s_k| < \infty$, which implies $\sum_{k=0}^{\infty} |d_k s_k| < \infty$. Therefore one has confirmed (6.1), indicating that we are justified in reversing the order of our generalized sums and the orthogonality relations under the above assumptions, which also are fulfilled for the polynomial families used throughout this paper.

In this section one has integral representations, infinite series, and representations in terms of the q -integral. In all the cases Lemma 6.1 can be applied and we are justified in interchanging the linear form and the infinite sum.

6.1 Definite integrals

6.1.1 The continuous q -ultraspherical/Rogers polynomials

The property of orthogonality for continuous q -ultraspherical/Rogers polynomials found in [10, (3.10.16)] is given by

$$\langle \mathbf{u}_R(\beta|q), C_m(x; \beta|q) C_n(x; \beta|q) \rangle = 2\pi \frac{(1-\beta)(\beta, q\beta; q)_\infty (\beta^2; q)_n}{(1-\beta q^n)(\beta^2, q; q)_\infty (q; q)_n} \delta_{mn}, \quad (6.2) \quad \boxed{\text{ortho}}$$

where the linear functional $\mathbf{u}_R(\beta|q)$ is defined in (3.4), and $w_R(x; \beta|q)$ is defined in (3.5). We will use this orthogonality relation for proofs of the following definite integrals.

Corollary 6.2. *Let $n \in \mathbb{N}_0$, $\beta, \gamma \in (-1, 1) \setminus \{0\}$, $0 < |q| < 1$, $|t| < 1 - \beta^2$. Then*

$$\begin{aligned} & \int_{-1}^1 (te^{-i\theta}; q)_\infty {}_2\phi_1 \left(\begin{matrix} \beta, \beta e^{2i\theta} \\ \beta^2 \end{matrix}; q, te^{-i\theta} \right) C_n(x; \gamma|q) \frac{w_R(x; \gamma|q)}{\sqrt{1-x^2}} dx \\ &= 2\pi (-\beta t)^n \frac{q^{\binom{n}{2}} (\gamma, q\gamma; q)_\infty (\beta, \gamma^2; q)_n}{(\gamma^2, q; q)_\infty (q, \beta^2, q\gamma; q)_n} {}_2\phi_5 \left(\begin{matrix} \beta\gamma^{-1}, \beta q^n \\ \gamma q^{n+1}, \pm\beta q^{n/2}, \pm\beta q^{(n+1)/2} \end{matrix}; q, \gamma(\beta t)^2 q^{2n+1} \right). \end{aligned}$$

Proof. Using the generalized generating function (3.2) and (6.2), the proof follows as above. ■

Corollary 6.3. *Let $n \in \mathbb{N}_0$, $\beta, \gamma \in (-1, 1) \setminus \{0\}$, $0 < |q| < 1$, $|t| < 1 - \beta^2$. Then*

$$\begin{aligned} & \int_{-1}^1 \frac{1}{(te^{i\theta}; q)_\infty} {}_2\phi_1 \left(\begin{matrix} \beta, \beta e^{2i\theta} \\ \beta^2 \end{matrix}; q, te^{-i\theta} \right) C_n(x; \gamma|q) \frac{w_R(x; \gamma|q)}{\sqrt{1-x^2}} dx \\ &= 2\pi t^n \frac{(\gamma, q\gamma; q)_\infty (\beta, \gamma^2; q)_n}{(\gamma^2, q; q)_\infty (q, \beta^2, q\gamma; q)_n} {}_6\phi_5 \left(\begin{matrix} \beta\gamma^{-1}, \beta q^n, 0, 0, 0, 0 \\ \gamma q^{n+1}, \pm\beta q^{n/2}, \pm\beta q^{(n+1)/2} \end{matrix}; q, \gamma t^2 \right). \end{aligned}$$

Proof. We complete the proof using (3.7) and (6.2). ■

Corollary 6.4. *Let $n \in \mathbb{N}_0$, $\gamma \in \mathbb{C}$, $\alpha, \beta \in (-1, 1) \setminus \{0\}$, $0 < |q| < 1$, $|t| < 1 - \beta^2$. Then*

$$\begin{aligned} & \int_{-1}^1 \frac{(\gamma te^{i\theta}; q)_\infty}{(te^{i\theta}; q)_\infty} {}_3\phi_2 \left(\begin{matrix} \gamma, \beta, \beta e^{2i\theta} \\ \beta^2, \gamma te^{i\theta} \end{matrix}; q, te^{-i\theta} \right) C_n(x; \alpha|q) \frac{w_R(x; \alpha|q)}{\sqrt{1-x^2}} dx \\ &= 2\pi t^n \frac{(\alpha, q\alpha; q)_\infty (\alpha^2, \gamma, \beta; q)_n}{(\alpha^2, q; q)_\infty (q, \beta^2, q\alpha; q)_n} {}_6\phi_5 \left(\begin{matrix} \beta/\alpha, \beta q^n, \pm(\gamma q^n)^{\frac{1}{2}} \pm (\gamma q^{n+1})^{\frac{1}{2}} \\ \alpha q^{n+1}, \pm\beta q^{n/2}, \pm\beta q^{(n+1)/2} \end{matrix}; q, \alpha t^2 \right). \end{aligned}$$

Proof. We complete the proof using (3.9) and (6.2). \blacksquare

$\langle \text{cor:22} \rangle?$ **Corollary 6.5.** Let $n \in \mathbb{N}_0$, $\beta, \gamma \in (-1, 1) \setminus \{0\}$, $0 < |q| < 1$, $|t| < \min\{(1 - \beta^2)(1 + \sqrt{q}|\beta|)(1 - q|\gamma|), 1\}$. Then

$$\begin{aligned} & \int_{-1}^1 {}_2\phi_1 \left(\begin{matrix} \pm\beta^{\frac{1}{2}}e^{i\theta} \\ -\beta \end{matrix}; q, te^{-i\theta} \right) {}_2\phi_1 \left(\begin{matrix} \pm(q\beta)^{\frac{1}{2}}e^{-i\theta} \\ -q\beta \end{matrix}; q, te^{i\theta} \right) C_n(x; \gamma|q) \frac{w_R(x; \gamma|q)}{\sqrt{1-x^2}} dx \\ &= 2\pi t^n \frac{(\gamma, q\gamma; q)_\infty (\gamma^2, \beta, \pm\beta q^{\frac{1}{2}}; q)_n}{(\gamma^2, q; q)_\infty (\beta^2, -q\beta, q\gamma, q; q)_n} \\ & \times {}_{10}\phi_9 \left(\begin{matrix} \beta\gamma^{-1}, \beta q^n, \pm(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}}, \pm i(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm i(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \\ \gamma q^{n+1}, \pm\beta q^{n/2}, \pm\beta q^{(n+1)/2}, \pm i(\beta q^{n+1})^{\frac{1}{2}}, \pm i(\beta q^{n+2})^{\frac{1}{2}} \end{matrix}; q, \gamma t^2 \right). \end{aligned}$$

Proof. We complete the proof using (3.11) and (6.2). \blacksquare

$\langle \text{cor:23} \rangle?$ **Corollary 6.6.** Let $n \in \mathbb{N}_0$, $\beta, \gamma \in (-1, 1) \setminus \{0\}$, $0 < |q| < 1$, $|t| < \min\{(1 - \beta^2)(1 + \sqrt{q}|\beta|)(1 - q|\gamma|), 1\}$. Then

$$\begin{aligned} & \int_{-1}^1 {}_2\phi_1 \left(\begin{matrix} \beta^{\frac{1}{2}}e^{i\theta}, (q\beta)^{\frac{1}{2}}e^{i\theta} \\ \beta q^{\frac{1}{2}} \end{matrix}; q, te^{-i\theta} \right) {}_2\phi_1 \left(\begin{matrix} -\beta^{\frac{1}{2}}e^{-i\theta}, -(q\beta)^{\frac{1}{2}}e^{-i\theta} \\ \beta q^{\frac{1}{2}} \end{matrix}; q, te^{i\theta} \right) \\ & \times C_n(x; \gamma|q) \frac{w_R(x; \gamma|q)}{\sqrt{1-x^2}} dx = 2\pi t^n \frac{(\gamma, q\gamma; q)_\infty (\gamma^2, \pm\beta, -\beta q^{\frac{1}{2}}; q)_n}{(\gamma^2, q; q)_\infty (\beta^2, \beta q^{\frac{1}{2}}, q\gamma, q; q)_n} \\ & \times {}_{10}\phi_9 \left(\begin{matrix} \beta\gamma^{-1}, \beta q^n, \pm i(\beta q^n)^{\frac{1}{2}}, \pm i(\beta q^{n+1})^{\frac{1}{2}}, \pm i(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}} \pm i(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \\ \gamma q^{n+1}, \pm\beta q^{n/2}, \pm\beta q^{(n+1)/2}, \pm(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \end{matrix}; q, \gamma t^2 \right). \end{aligned}$$

Proof. We complete the proof using (3.13) and (6.2). \blacksquare

$\langle \text{cor:24} \rangle?$ **Corollary 6.7.** Let $n \in \mathbb{N}_0$, $\beta, \gamma \in (-1, 1) \setminus \{0\}$, $0 < |q| < 1$, $|t| < \min\{(1 - \beta^2)(1 + \sqrt{q}|\beta|), 1\}$. Then

$$\begin{aligned} & \int_{-1}^1 {}_2\phi_1 \left(\begin{matrix} \beta^{\frac{1}{2}}e^{i\theta}, -(q\beta)^{\frac{1}{2}}e^{i\theta} \\ -\beta q^{\frac{1}{2}} \end{matrix}; q, te^{-i\theta} \right) {}_2\phi_1 \left(\begin{matrix} (q\beta)^{\frac{1}{2}}e^{-i\theta}, -\beta^{\frac{1}{2}}e^{-i\theta} \\ -\beta q^{\frac{1}{2}} \end{matrix}; q, te^{i\theta} \right) \\ & \times C_n(x; \gamma|q) \frac{w_R(x; \gamma|q)}{\sqrt{1-x^2}} dx = 2\pi t^n \frac{(\gamma, q\gamma; q)_\infty (\gamma^2, \pm\beta\beta q^{\frac{1}{2}}; q)_n}{(\gamma^2, q; q)_\infty (\beta^2, -\beta q^{\frac{1}{2}}, q\gamma, q; q)_n} \\ & \times {}_{10}\phi_9 \left(\begin{matrix} \beta\gamma^{-1}, \beta q^n, \pm i(\beta q^n)^{\frac{1}{2}}, \pm i(\beta q^{n+1})^{\frac{1}{2}}, \pm(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \\ \gamma q^{n+1}, \pm\beta q^{n/2}, \pm\beta q^{(n+1)/2}, \pm i(\beta q^{n+\frac{1}{2}})^{\frac{1}{2}}, \pm i(\beta q^{n+\frac{3}{2}})^{\frac{1}{2}} \end{matrix}; q, \gamma t^2 \right). \end{aligned}$$

Proof. We complete the proof using (3.14) and (6.2). \blacksquare

6.1.2 The q -Laguerre polynomials

The continuous orthogonality relation for the q -Laguerre polynomials is given by the following result. Notice that this result appears in [2, Section 2].

$\langle \text{theo:25} \rangle?$ **Proposition 6.8.** Let $\alpha \in (-1, \infty)$, $m, n \in \mathbb{N}_0$, $0 < |q| < 1$. Then

$$\int_0^\infty L_m^{(\alpha)}(x; q) L_n^{(\alpha)}(x; q) \frac{x^\alpha}{(-x; q)_\infty} dx = -\frac{\delta_{m,n}}{q^n} \begin{cases} \frac{\pi(q^{-\alpha}; q)_\infty (q^{\alpha+1}; q)_n}{\sin(\pi\alpha)(q; q)_\infty (q; q)_n}, & \text{if } \alpha \in (-1, \infty) \setminus \mathbb{N}_0, \\ \frac{(q^{n+1}; q)_\alpha \log q}{q^{\alpha(\alpha+1)/2}}, & \text{if } \alpha \in \mathbb{N}_0. \end{cases} \quad (6.3) \quad \boxed{\text{HSC1}}$$

Proof. The continuous orthogonality relation for the q -Laguerre polynomials is given in [10, (14.21.2)] with the right-hand side expressed in terms of gamma functions, namely

$$\int_0^\infty L_m^{(\alpha)}(x; q) L_n^{(\alpha)}(x; q) \frac{x^\alpha}{(-x; q)_\infty} dx = \frac{(q^{-\alpha}; q)_\infty (q^{\alpha+1}; q)_n}{q^n (q; q)_\infty (q; q)_n} \Gamma(-\alpha) \Gamma(\alpha + 1) \delta_{m,n}.$$

The gamma functions can be replaced using the reflection formula [7, (5.5.3)] and the result is given in the theorem for $\alpha \in (-1, \infty) \setminus \mathbb{N}_0$. The result for $\alpha \in \mathbb{N}_0$ is a consequence of (2.3) and [1, cf. (2.9)], namely

$$\lim_{\alpha \rightarrow k} \frac{(q^{1-\alpha}; q)_\infty}{\sin(\pi\alpha) (aq^{-\alpha}; q)_\infty} = \frac{-(q; q)_\infty (q; q)_{k-1} \log q}{\pi q^{\binom{k}{2}} (a; q)_\infty (a; q^{-1})_k},$$

which leads to

$$\lim_{\alpha \rightarrow k} \frac{(q^{-\alpha}; q)_\infty}{\sin(\pi\alpha)} = \frac{(q; q)_\infty (q; q)_k \log q}{\pi q^{k(k+1)/2}}.$$

Applying this limit completes the proof. ■

^{?(cor:26)?} **Corollary 6.9.** Let $n \in \mathbb{N}_0$, $\alpha, \beta \in (-1, \infty)$, $0 < |q| < 1$, $|t| < (1 - q^{\alpha+1})(1 - q)$. Then

$$\begin{aligned} & \int_0^\infty {}_0\phi_1 \left(\begin{matrix} - \\ q^{\alpha+1} \end{matrix}; q, -xtq^{\alpha+1} \right) L_n^{(\beta)}(x; q) \frac{x^\beta}{(-x; q)_\infty} dx \\ &= \frac{-(tq^{\alpha-\beta})^n (t; q)_\infty}{q^n (q^{\alpha+1}; q)_n} {}_2\phi_1 \left(\begin{matrix} q^{\alpha-\beta}, 0 \\ q^{\alpha+n+1} \end{matrix}; q, t \right) \begin{cases} \frac{\pi (q^{-\beta}; q)_\infty (q^{\beta+1}; q)_n}{\sin(\pi\beta) (q; q)_\infty (q; q)_n}, & \text{if } \beta \in (-1, \infty) \setminus \mathbb{N}_0, \\ \frac{(q^{n+1}; q)_\beta \log q}{q^{\beta(\beta+1)/2}}, & \text{if } \beta \in \mathbb{N}_0. \end{cases} \end{aligned}$$

Proof. Using (6.3) with (5.2) completes the proof. ■

^{?(cor:27)?} **Corollary 6.10.** Let $n \in \mathbb{N}_0$, $\alpha, \beta \in (-1, \infty)$, $0 < |q| < 1$, $|t| < (1 - q^{\alpha+1})(1 - q)$. Then

$$\begin{aligned} & \int_0^\infty {}_0\phi_2 \left(\begin{matrix} - \\ q^{\alpha+1}, t \end{matrix}; q, -xtq^{\alpha+1} \right) L_n^{(\beta)}(x; q) \frac{x^\beta}{(-x; q)_\infty} dx \\ &= \frac{-(-tq^{\alpha-\beta})^n}{q^n (t; q)_\infty (q^{\alpha+1}; q)_n} {}_1\phi_1 \left(\begin{matrix} q^{\alpha-\beta} \\ q^{\alpha+n+1} \end{matrix}; q, tq^n \right) \begin{cases} \frac{\pi (q^{-\beta}; q)_\infty (q^{\beta+1}; q)_n}{\sin(\pi\beta) (q; q)_\infty (q; q)_n}, & \text{if } \beta \in (-1, \infty) \setminus \mathbb{N}_0, \\ \frac{(q^{n+1}; q)_\beta \log q}{q^{\beta(\beta+1)/2}}, & \text{if } \beta \in \mathbb{N}_0. \end{cases} \end{aligned}$$

Proof. Using (6.3) with (5.5) completes the proof. ■

^{?(cor:28)?} **Corollary 6.11.** Let $n \in \mathbb{N}_0$, $\alpha, \beta \in (-1, \infty)$, $\gamma \in \mathbb{C}$, $0 < |q| < 1$, $|t| < 1 - q$. Then

$$\begin{aligned} & \int_0^\infty {}_1\phi_2 \left(\begin{matrix} \gamma \\ q^{\alpha+1}, \gamma t \end{matrix}; q, -xtq^{\alpha+1} \right) L_n^{(\beta)}(x; q) \frac{x^\beta}{(-x; q)_\infty} dx = \frac{-(tq^{\alpha-\beta})^n (t; q)_\infty (\gamma; q)_n}{q^n (\gamma t; q)_\infty (q^{\alpha+1}; q)_n} \\ & \quad \times {}_2\phi_1 \left(\begin{matrix} q^{\alpha-\beta}, \gamma q^n \\ q^{\alpha+n+1} \end{matrix}; q, t \right) \begin{cases} \frac{\pi (q^{-\beta}; q)_\infty (q^{\beta+1}; q)_n}{\sin(\pi\beta) (q; q)_\infty (q; q)_n}, & \text{if } \beta \in (-1, \infty) \setminus \mathbb{N}_0, \\ \frac{(q^{n+1}; q)_\beta \log q}{q^{\beta(\beta+1)/2}}, & \text{if } \beta \in \mathbb{N}_0. \end{cases} \end{aligned}$$

Proof. Using (6.3) with (5.7) completes the proof. ■

6.2 Infinite series

6.2.1 The little q -Laguerre/Wall polynomials

The little q -Laguerre/Wall polynomials satisfy a discrete orthogonality relation, namely [10, (14.20.2)]

$$\sum_{k=0}^{\infty} p_m(q^k; a|q) p_n(q^k; a|q) \frac{(aq)^k}{(q; q)_k} = \frac{(aq)^n (q; q)_n}{(aq; q)_{\infty} (aq; q)_n} \delta_{m,n},$$

for $a \in (0, 1/q)$, with $0 < |q| < 1$.

^{?(cor:29)?} **Corollary 6.12.** *Let $n \in \mathbb{N}_0$, $0 < |q| < 1$, $\alpha, \beta \in (0, q^{-1})$, $|t| < \min\{(1 - \beta^2)(1 + \sqrt{q}|\beta|), 1\}$. Then*

$$\sum_{k=0}^{\infty} \frac{(q\beta)^k}{(tq^k; q)_{\infty}} {}_0\phi_1\left(\begin{matrix} - \\ q\alpha \end{matrix}; q, tq^k\right) \frac{p_n(q^k; \beta|q)}{(q; q)_k} = \frac{q^{\binom{n}{2}} (-q\beta t)^n}{(t, q\beta; q)_{\infty} (q\alpha; q)_n} {}_1\phi_1\left(\begin{matrix} \alpha/\beta \\ \alpha q^{n+1} \end{matrix}; q, t\beta q^{n+1}\right).$$

Proof. We begin with the generalized generating function (3.14) and using (6.2) completes the proof. This orthogonality isn't there. ■

6.2.2 The q -Laguerre polynomials

One type of discrete orthogonality that the q -Laguerre polynomials satisfy is [10, (14.21.3)]

$$\sum_{k=-\infty}^{\infty} L_m^{(\alpha)}(cq^k; q) L_n^{(\alpha)}(cq^k; q) \frac{q^{(\alpha+1)k}}{(-cq^k; q)_{\infty}} = \frac{(q, -cq^{\alpha+1}, -q^{-\alpha}/c; q)_{\infty} (q^{\alpha+1}; q)_n}{q^n (q^{\alpha+1}, -c, -q/c; q)_{\infty} (q; q)_n} \delta_{m,n}, \quad (6.4) \quad \boxed{\text{bilateralorthogqL}}$$

for $\alpha \in (-1, \infty)$, $c > 0$.

^{?(cor:30)?} **Corollary 6.13.** *Let $n \in \mathbb{N}_0$, $0 < |q| < 1$, $\alpha, \beta \in (-1, \infty)$, $|t| < (1 - q^{\alpha+1})(1 - q)$, $c > 0$. Then*

$$\begin{aligned} \sum_{k=-\infty}^{\infty} {}_0\phi_1\left(\begin{matrix} - \\ q^{\alpha+1} \end{matrix}; q, -ctq^{\alpha+k+1}\right) L_n^{(\beta)}(cq^k; q) \frac{q^{(\beta+1)k}}{(-cq^k; q)_{\infty}} \\ = \frac{(tq^{\alpha-\beta})^n (t, q, -cq^{\beta+1}, -q^{-\beta}/c; q)_{\infty} (q^{\beta+1}; q)_n}{q^n (q^{\beta+1}, -c, -q/c; q)_{\infty} (q, q^{\alpha+1}; q)_n} {}_2\phi_1\left(\begin{matrix} q^{\alpha-\beta}, 0 \\ q^{\alpha+n+1} \end{matrix}; q, t\right). \end{aligned}$$

Proof. This follows using (5.2) with (6.4). ■

^{?(cor:31)?} **Corollary 6.14.** *Let $n \in \mathbb{N}_0$, $0 < |q| < 1$, $\alpha, \beta \in (-1, \infty)$, $|t| < (1 - q^{\alpha+1})(1 - q)$, $c > 0$. Then*

$$\begin{aligned} \sum_{k=-\infty}^{\infty} {}_0\phi_2\left(\begin{matrix} - \\ q^{\alpha+1}, t \end{matrix}; q, -ctq^{\alpha+k+1}\right) L_n^{(\beta)}(cq^k; q) \frac{q^{(\beta+1)k}}{(-cq^k; q)_{\infty}} \\ = \frac{(-tq^{\alpha-\beta})^n q^{\binom{n}{2}} (q, -cq^{\beta+1}, -q^{-\beta}/c; q)_{\infty} (q^{\beta+1}; q)_n}{q^n (t, q^{\beta+1}, -c, -q/c; q)_{\infty} (q, q^{\alpha+1}; q)_n} {}_1\phi_1\left(\begin{matrix} q^{\alpha-\beta} \\ q^{\alpha+n+1} \end{matrix}; q, tq^n\right). \end{aligned}$$

Proof. This follows using (5.5) with (6.4). ■

^{?(cor:32)?} **Corollary 6.15.** *Let $n \in \mathbb{N}_0$, $0 < |q| < 1$, $\alpha, \beta \in (-1, \infty)$, $\gamma \in \mathbb{C}$, $|t| < 1 - q$, $c > 0$. Then*

$$\begin{aligned} \sum_{k=-\infty}^{\infty} {}_1\phi_2\left(\begin{matrix} \gamma \\ q^{\alpha+1}, \gamma t \end{matrix}; q, -ctq^{\alpha+k+1}\right) L_n^{(\beta)}(cq^k; q) \frac{q^{(\beta+1)k}}{(-cq^k; q)_{\infty}} \\ = \frac{(tq^{\alpha-\beta})^n (t, q, -cq^{\beta+1}, -q^{-\beta}/c; q)_{\infty} (\gamma, q^{\beta+1}; q)_n}{q^n (\gamma t, q^{\beta+1}, -c, -q/c; q)_{\infty} (q, q^{\alpha+1}; q)_n} {}_2\phi_1\left(\begin{matrix} q^{\alpha-\beta}, \gamma q^n \\ q^{\alpha+n+1} \end{matrix}; q, t\right). \end{aligned}$$

Proof. This follows using (5.7) with (6.4). ■

6.3 The q -Integrals

6.3.1 The q -Laguerre polynomials

One type of orthogonality for the q -Laguerre polynomials is [10, (14.21.4)]

$$\int_0^\infty L_m^{(\alpha)}(x; q) L_n^{(\alpha)}(x; q) \frac{x^\alpha}{(-x; q)_\infty} d_q x = \frac{(1-q)(q, -q^{\alpha+1}, -q^{-\alpha}; q)_\infty (q^{\alpha+1}; q)_n}{2q^n (q^{\alpha+1}, -q, -q; q)_\infty (q; q)_n} \delta_{m,n}. \quad (6.5) \quad \boxed{\text{Jackson orthog } q\text{-Lag}}$$

Using this orthogonality relation we can obtain new q -integrals using our generalized generating functions for the q -Laguerre polynomials.

^{?(cor:33)?} **Corollary 6.16.** *Let $n \in \mathbb{N}_0$, $0 < |q| < 1$, $\alpha, \beta \in (-1, \infty)$, $|t| < (1 - q^{\alpha+1})(1 - q)$. Then*

$$\begin{aligned} \int_0^\infty {}_0\phi_1\left(\begin{matrix} - \\ q^{\alpha+1} \end{matrix}; q, -xtq^{\alpha+1}\right) L_n^{(\beta)}(x; q) \frac{x^\beta}{(-x; q)_\infty} d_q x \\ = \frac{(1-q)(tq^{\alpha-\beta})^n (t, q, -q^{\beta+1}, -q^{-\beta}; q)_\infty (q^{\beta+1}; q)_n}{2q^n (q^{\beta+1}, -q, -q; q)_\infty (q, q^{\alpha+1}; q)_n} {}_2\phi_1\left(\begin{matrix} q^{\alpha-\beta}, 0 \\ q^{\alpha+n+1} \end{matrix}; q, t\right). \end{aligned}$$

Proof. Using (5.2) with (6.5) completes this proof. ■

^{?(cor:34)?} **Corollary 6.17.** *Let $n \in \mathbb{N}_0$, $0 < |q| < 1$, $\alpha, \beta \in (-1, \infty)$, $|t| < (1 - q^{\alpha+1})(1 - q)$. Then*

$$\begin{aligned} \int_0^\infty {}_0\phi_2\left(\begin{matrix} - \\ q^{\alpha+1}, t \end{matrix}; q, -xtq^{\alpha+1}\right) L_n^{(\beta)}(x; q) \frac{x^\beta}{(-x; q)_\infty} d_q x \\ = \frac{(1-q)(-tq^{\alpha-\beta})^n q^{\binom{n}{2}} (q, -q^{\beta+1}, -q^{-\beta}; q)_\infty (q^{\beta+1}; q)_n}{2q^n (t, q^{\beta+1}, -q, -q; q)_\infty (q, q^{\alpha+1}; q)_n} {}_1\phi_1\left(\begin{matrix} q^{\alpha-\beta} \\ q^{\alpha+n+1} \end{matrix}; q, tq^n\right). \end{aligned}$$

Proof. Using (5.5) with (6.5) completes this proof. ■

^{?(cor:35)?} **Corollary 6.18.** *Let $n \in \mathbb{N}_0$, $0 < |q| < 1$, $\alpha, \beta \in (-1, \infty)$, $\gamma \in \mathbb{C}$, $|t| < 1 - q$. Then*

$$\begin{aligned} \int_0^\infty {}_1\phi_2\left(\begin{matrix} \gamma \\ q^{\alpha+1}, \gamma t \end{matrix}; q, -xtq^{\alpha+1}\right) L_n^{(\beta)}(x; q) \frac{x^\beta}{(-x; q)_\infty} d_q x \\ = \frac{(1-q)(tq^{\alpha-\beta})^n (t, q, -q^{\beta+1}, -q^{-\beta}; q)_\infty (\gamma, q^{\beta+1}; q)_n}{2q^n (\gamma t, q^{\beta+1}, -q, -q; q)_\infty (q, q^{\alpha+1}; q)_n} {}_2\phi_1\left(\begin{matrix} q^{\alpha-\beta}, \gamma q^n \\ q^{\alpha+n+1} \end{matrix}; q, t\right). \end{aligned}$$

Proof. Using (5.7) with (6.5) completes this proof. ■

Acknowledgements

Much thanks to Hans Volkmer for valuable discussions. The author R. S. Costas-Santos acknowledges financial support by Dirección General de Investigación, Ministerio de Economía y Competitividad of Spain, grant MTM2015-65888-C4-2-P.

References

- Askey80 [1] R. Askey. Ramanujan’s extensions of the gamma and beta functions. *The American Mathematical Monthly*, 87(5):346–359, 1980.
- Christiansen [2] J. S. Christiansen. The moment problem associated with the q -Laguerre polynomials. *Constructive Approximation. An International Journal for Approximations and Expansions*, 19(1):1–22, 2003.

- [3] [Cohl12pow](#) H. S. Cohl. Fourier, Gegenbauer and Jacobi expansions for a power-law fundamental solution of the polyharmonic equation and polyspherical addition theorems. *Symmetry, Integrability and Geometry: Methods and Applications*, 9(042):26, 2013.
- [4] [CohlGenGegen](#) H. S. Cohl. On a generalization of the generating function for Gegenbauer polynomials. *Integral Transforms and Special Functions*, 24(10):807–816, 2013.
- [5] [CostasWakhare2019](#) H. S. Cohl, R. S. Costas-Santos, and T. V. Wakhare. On a generalization of the Rogers generating function. *Journal of Mathematical Analysis and Applications*, 475(2):1019–1043, 2019.
- [6] [CohlMacKVolk](#) H. S. Cohl, C. MacKenzie, and H. Volkmer. Generalizations of generating functions for hypergeometric orthogonal polynomials with definite integrals. *Journal of Mathematical Analysis and Applications*, 407(2):211–225, 2013.
- [7] [NIST:DLMF](#) *NIST Digital Library of Mathematical Functions*. <http://dlmf.nist.gov>, Release 1.0.27 of 2020-06-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [8] [GaspRah](#) G. Gasper and M. Rahman. *Basic hypergeometric series*, volume 96 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 2004. With a foreword by Richard Askey.
- [9] [Ismail:2009:CQ0](#) M. E. H. Ismail. *Classical and Quantum Orthogonal Polynomials in One Variable*, volume 98 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2009. With two chapters by Walter Van Assche, With a foreword by Richard A. Askey, Corrected reprint of the 2005 original.
- [10] [Koekoeketal](#) R. Koekoek, P. A. Lesky, and R. F. Swarttouw. *Hypergeometric orthogonal polynomials and their q -analogues*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2010. With a foreword by Tom H. Koornwinder.